

**Jemperli**   
(dostarlimab-gxly) Injection 500 mg

# COUNSELING PATIENTS

## Who Are Receiving JEMPERLI



Not an actual patient  
taking JEMPERLI.

### Indications

- JEMPERLI, in combination with carboplatin and paclitaxel, followed by JEMPERLI as a single agent, is indicated for the treatment of adult patients with primary advanced or recurrent endometrial cancer (EC).
- JEMPERLI, as a single agent, is indicated for the treatment of adult patients with mismatch repair deficient (dMMR) recurrent or advanced EC, as determined by an FDA-approved test, that has progressed on or following prior treatment with a platinum-containing regimen in any setting and are not candidates for curative surgery or radiation.

### Important Safety Information

#### Severe and Fatal Immune-Mediated Adverse Reactions

- Immune-mediated adverse reactions, which can be severe or fatal, can occur in any organ system or tissue and can occur at any time during or after treatment with a PD-1/PD-L1-blocking antibody, including JEMPERLI.
- Monitor closely for signs and symptoms of immune-mediated adverse reactions. Evaluate liver enzymes, creatinine, and thyroid function tests at baseline and periodically during treatment. For suspected immune-mediated adverse reactions, initiate appropriate workup to exclude alternative etiologies, including infection. Institute medical management promptly, including specialty consultation as appropriate.
- Based on the severity of the adverse reaction, withhold or permanently discontinue JEMPERLI. In general, if JEMPERLI requires interruption or discontinuation, administer systemic corticosteroids (1 to 2 mg/kg/day prednisone or equivalent) until improvement to  $\leq$ Grade 1. Upon improvement to  $\leq$ Grade 1, initiate corticosteroid taper and continue to taper over at least 1 month. Consider administration of other systemic immunosuppressants in patients whose immune-mediated adverse reaction is not controlled with corticosteroids.

FDA, US Food and Drug Administration; PD-1, programmed death receptor-1; PD-L1, programmed death ligand-1.

**Please see additional Important Safety Information on the next pages and full [Prescribing Information](#), including [Medication Guide](#) for patients.**

# Inform patients of the risk of immune-mediated adverse reactions that may be severe or fatal, may occur after discontinuation of treatment, and may require corticosteroid or other treatment and interruption or discontinuation of JEMPERLI.

## Immune-mediated adverse reactions may include:



### Pneumonitis

Advise patients to contact their healthcare provider immediately for new or worsening cough, chest pain, or shortness of breath



### Colitis

Advise patients to contact their healthcare provider immediately for diarrhea or severe abdominal pain



### Hepatitis

Advise patients to contact their healthcare provider immediately for jaundice, severe nausea or vomiting, or easy bruising or bleeding



### Immune-mediated endocrinopathies

Advise patients to contact their healthcare provider immediately for signs or symptoms of hypothyroidism, hyperthyroidism, thyroiditis, adrenal insufficiency, hypophysitis, or type 1 diabetes mellitus



### Nephritis

Advise patients to contact their healthcare provider immediately for signs or symptoms of nephritis



### Severe skin reactions

Advise patients to contact their healthcare provider immediately for any signs or symptoms of severe skin reactions, SJS, TEN, or DRESS



### Other immune-mediated adverse reactions

- Advise patients that immune-mediated adverse reactions can occur and may involve any organ system, and to contact their healthcare provider immediately for any new signs or symptoms
- Advise patients of the risk of solid organ transplant rejection and to contact their healthcare provider immediately for signs or symptoms of organ transplant rejection

## Other information to consider:

### • Infusion-related reactions

Advise patients to contact their healthcare provider immediately for signs or symptoms of infusion-related reactions

### • Complications of allogeneic HSCT

Advise patients of the risk of post-allogeneic hematopoietic stem cell transplantation (HSCT) complications

### • Embryo-fetal toxicity

- Advise females of reproductive potential of the potential risk to a fetus and to inform their healthcare provider of a known or suspected pregnancy
- Advise females of reproductive potential to use effective contraception during treatment with JEMPERLI and for 4 months after the last dose

### • Lactation

Advise women not to breastfeed during treatment with JEMPERLI and for 4 months after the last dose

Healthcare providers should refer to the Patient Counseling Information in Section 17 when counseling patients and advise patients to read the [Medication Guide](#) in the [Prescribing Information](#).

## Important Safety Information (continued)

### Immune-Mediated Pneumonitis

- JEMPERLI can cause immune-mediated pneumonitis, which can be fatal. In patients treated with other PD-1/PD-L1-blocking antibodies, the incidence of pneumonitis is higher in patients who have received prior thoracic radiation. Pneumonitis occurred in 2.3% (14/605) of patients, including Grade 2 (1.3%), Grade 3 (0.8%), and Grade 4 (0.2%) pneumonitis.

### Immune-Mediated Colitis

- Colitis occurred in 1.3% (8/605) of patients, including Grade 2 (0.7%) and Grade 3 (0.7%) adverse reactions. Cytomegalovirus infection/reactivation have occurred in patients with corticosteroid-refractory immune-mediated colitis. In such cases, consider repeating infectious workup to exclude alternative etiologies.

## Important Safety Information (continued)

### Immune-Mediated Hepatitis

- JEMPERLI can cause immune-mediated hepatitis, which can be fatal. Grade 3 hepatitis occurred in 0.5% (3/605) of patients.

### Immune-Mediated Endocrinopathies

- Adrenal Insufficiency
  - Adrenal insufficiency occurred in 1.2% (7/605) of patients, including Grade 2 (0.5%) and Grade 3 (0.7%). For Grade 2 or higher adrenal insufficiency, initiate symptomatic treatment per institutional guidelines, including hormone replacement as clinically indicated. Withhold or permanently discontinue JEMPERLI depending on severity.

## Important Safety Information (continued)

### Immune-Mediated Endocrinopathies (continued)

- Hypophysitis
  - JEMPERLI can cause immune-mediated hypophysitis. Grade 3 hypophysitis occurred in 0.4% (1/241) of patients receiving JEMPERLI in combination with carboplatin and paclitaxel. Grade 2 hypophysitis occurred in 0.2% (1/605) of patients receiving JEMPERLI as a single agent. Initiate hormone replacement as clinically indicated. Withhold or permanently discontinue JEMPERLI depending on severity.

**DRESS**, drug rash with eosinophilia and systemic symptoms; **PD-1**, programmed death receptor-1; **PD-L1**, programmed death ligand-1; **SJS**, Stevens-Johnson syndrome; **TEN**, toxic epidermal necrolysis.

Please see remaining Important Safety Information on the next page and full [Prescribing Information](#), including [Medication Guide](#) for patients.

## Important Safety Information (continued)

### Immune-Mediated Endocrinopathies (continued)

- Thyroid Disorders
  - Grade 2 thyroiditis occurred in 0.5% (3/605) of patients. Grade 2 hypothyroidism occurred in 12% (30/241) of patients receiving JEMPERLI in combination with carboplatin and paclitaxel. Grade 2 hypothyroidism occurred in 8% (46/605) of patients receiving JEMPERLI as a single agent. Hyperthyroidism occurred in 3.3% (8/241) of patients receiving JEMPERLI in combination with carboplatin and paclitaxel, including Grade 2 (2.9%) and Grade 3 (0.4%). Hyperthyroidism occurred in 2.3% (14/605) of patients receiving JEMPERLI as a single agent, including Grade 2 (2.1%) and Grade 3 (0.2%). Initiate thyroid hormone replacement or medical management of hyperthyroidism as clinically indicated. Withhold or permanently discontinue JEMPERLI depending on severity.
- Type 1 Diabetes Mellitus, Which Can Present with Diabetic Ketoacidosis
  - JEMPERLI can cause type 1 diabetes mellitus, which can present with diabetic ketoacidosis. Grade 3 type 1 diabetes mellitus occurred in 0.4% (1/241) of patients receiving JEMPERLI in combination with carboplatin and paclitaxel. Grade 3 type 1 diabetes mellitus occurred in 0.2% (1/605) of patients receiving JEMPERLI as a single agent. Monitor patients for hyperglycemia or other signs and symptoms of diabetes. Initiate treatment with insulin as clinically indicated. Withhold or permanently discontinue JEMPERLI depending on severity.

### Immune-Mediated Nephritis with Renal Dysfunction

- JEMPERLI can cause immune-mediated nephritis, which can be fatal. Grade 2 nephritis, including tubulointerstitial nephritis, occurred in 0.5% (3/605) of patients.

### Immune-Mediated Dermatologic Adverse Reactions

- JEMPERLI can cause immune-mediated rash or dermatitis. Bullous and exfoliative dermatitis, including Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN), and drug rash with eosinophilia and systemic symptoms (DRESS), have occurred with PD-1/PD-L1–blocking antibodies. Topical emollients and/or topical corticosteroids may be adequate to treat mild to moderate non-bullous/exfoliative rashes. Withhold or permanently discontinue JEMPERLI depending on severity.

## Important Safety Information (continued)

### Other Immune-Mediated Adverse Reactions

- The following clinically significant immune-mediated adverse reactions occurred in <1% of the 605 patients treated with JEMPERLI or were reported with the use of other PD-1/PD-L1–blocking antibodies. Severe or fatal cases have been reported for some of these adverse reactions.
  - *Nervous System*: Meningitis, encephalitis, myelitis and demyelination, myasthenic syndrome/myasthenia gravis, Guillain-Barré syndrome, nerve paresis, autoimmune neuropathy
  - *Cardiac/Vascular*: Myocarditis, pericarditis, vasculitis
  - *Ocular*: Uveitis, iritis, other ocular inflammatory toxicities. Some cases can be associated with retinal detachment. Various grades of visual impairment to include blindness can occur
  - *Gastrointestinal*: Pancreatitis, including increases in serum amylase and lipase levels, gastritis, duodenitis
  - *Musculoskeletal and Connective Tissue*: Myositis/polymyositis, rhabdomyolysis and associated sequelae including renal failure, arthritis, polymyalgia rheumatica
  - *Endocrine*: Hypoparathyroidism
  - *Other (Hematologic/Immune)*: Autoimmune hemolytic anemia, aplastic anemia, hemophagocytic lymphohistiocytosis, systemic inflammatory response syndrome, histiocytic necrotizing lymphadenitis (Kikuchi lymphadenitis), sarcoidosis, immune thrombocytopenia, solid organ transplant rejection, other transplant (including corneal graft) rejection

### Infusion-Related Reactions

- Severe or life-threatening infusion-related reactions have been reported with PD-1/PD-L1–blocking antibodies. Severe infusion-related reactions (Grade 3) occurred in 0.2% (1/605) of patients receiving JEMPERLI. Monitor patients for signs and symptoms of infusion-related reactions. Interrupt or slow the rate of infusion or permanently discontinue JEMPERLI based on severity of reaction.

### Complications of Allogeneic HSCT

- Fatal and other serious complications can occur in patients who receive allogeneic hematopoietic stem cell transplantation (HSCT) before or after treatment with a PD-1/PD-L1–blocking antibody, which may occur despite intervening therapy. Monitor patients closely for transplant-related complications and intervene promptly.

## Important Safety Information (continued)

### Embryo-Fetal Toxicity and Lactation

- Based on its mechanism of action, JEMPERLI can cause fetal harm. Advise pregnant women of the potential risk to a fetus. Advise females of reproductive potential to use effective contraception during treatment with JEMPERLI and for 4 months after their last dose. Because of the potential for serious adverse reactions from JEMPERLI in a breastfed child, advise women not to breastfeed during treatment with JEMPERLI and for 4 months after their last dose.

### Common Adverse Reactions

The most common adverse reactions (≥20%), including laboratory abnormalities, in patients with EC who received JEMPERLI in combination with carboplatin and paclitaxel were decreased hemoglobin, increased creatinine, peripheral neuropathy, decreased white blood cell count, fatigue, nausea, alopecia, decreased platelets, increased glucose, decreased lymphocytes, decreased magnesium, decreased neutrophils, increased AST, arthralgia, rash, constipation, diarrhea, increased ALT, decreased potassium, decreased albumin, decreased sodium, increased alkaline phosphatase, abdominal pain, dyspnea, decreased appetite, increased amylase, decreased phosphate, urinary tract infection, and vomiting.

The most common adverse reactions (≥20%) in patients with dMMR EC who received JEMPERLI as a single agent were fatigue/asthenia, anemia, nausea, diarrhea, constipation, vomiting, and rash. The most common Grade 3 or 4 laboratory abnormalities (>2%) were decreased lymphocytes, decreased sodium, increased alanine aminotransferase, increased creatinine, decreased neutrophils, decreased albumin, and increased alkaline phosphatase.

Please see full [Prescribing Information, including Medication Guide for patients](#).

For more information, please visit [www.jemperlihcp.com](http://www.jemperlihcp.com)

ALT, alanine aminotransferase; AST, aspartate aminotransferase; dMMR, mismatch repair deficient; EC, endometrial cancer; PD-1, programmed death receptor-1; PD-L1, programmed death ligand-1.

Reference: JEMPERLI. Prescribing Information. GSK; 2024.

## Together with GSK Oncology

One source for GSK access and reimbursement services



Co-pay assistance program for commercial patients



Coverage support



Claims assistance



Patient Assistance Program (PAP) for uninsured and Medicare patients



Referral to third-party support services

Phone: 1-844-447-5662

Online Application and Resources: [www.TogetherwithGSKOncology.com](http://www.TogetherwithGSKOncology.com)

Together with GSK Oncology provides resources for patients and healthcare professionals. Specific eligibility requirements are determined by the payer; therefore, patients and healthcare professionals should confirm information directly with payers. Together with GSK Oncology does not guarantee coverage or payer reimbursement.

Trademarks are owned by or licensed to the GSK group of companies.



©2024 GSK or licensor.  
PMUS-DSTLBND240032 September 2024  
Produced in USA.

File Name: Patient Counseling Guide 2024

**Jemperli**  
(dostarlimab-gxly) Injection 500 mg